



Organic makes a difference

the nutritional benefits of organically produced foods

Marcin Barański, Chris Seal, Roy Sanderson, Gavin Stewart and Carlo Leifert

Nafferton Ecological Farming Group (NEFG)
School of Agriculture, Food and Rural Development
Newcastle University



New evidence for significant and nutritionally relevant composition differences



British Journal of Nutrition (2014), **112**, 794–811

doi:10.1017/S0007114514001366

© The Authors 2014. The online version of this article is published within an Open Access environment subject to the conditions of the Creative Commons Attribution licence <http://creativecommons.org/licenses/by/3.0/>

Higher antioxidant and lower cadmium concentrations and lower incidence of pesticide residues in organically grown crops: a systematic literature review and meta-analyses

Marcin Barański¹, Dominika Średnicka-Tober¹, Nikolaos Volakakis¹, Chris Seal², Roy Sanderson³, Gavin B. Stewart¹, Charles Benbrook⁴, Bruno Biavati⁵, Emilia Markellou⁶, Charilaos Giotis⁷, Joanna Gromadzka-Ostrowska⁸, Ewa Rembiałkowska⁸, Krystyna Skwarło-Sońta⁹, Raija Tahvonen¹⁰, Dagmar Janovská¹¹, Urs Niggli¹², Philippe Nicot¹³ and Carlo Leifert^{1*}

¹*School of Agriculture, Food and Rural Development, Newcastle University, Nafferton Farm, Stocksfield, Northumberland, NE43 7XD, UK*

²*Human Nutrition Research Centre, School of Agriculture, Food and Rural Development, Newcastle University, Agriculture Building, Kings Road, Newcastle upon Tyne NE1 7RU, UK*

³*School of Biology, Newcastle University, Ridley Building, Newcastle upon Tyne NE1 7RU, UK*

⁴*Center for Sustaining Agriculture and Natural Resources, Washington State University, Pullman, WA, USA*

⁵*Department of Agricultural Sciences, School of Agriculture and Veterinary Medicine, University of Bologna, Viale Fanin 42, 40127 Bologna, Italy*

⁶*Department of Pesticide Control and*


⁷*Department of Organic Farming and*

⁸*Ioanif Momferatou & Ilia Miniati PC 28100, Argostoli, Cephalonia, Greece*

⁹*Department of Human Nutrition and Consumer Sciences, Wageningen University of Life Sciences, Wageningen, 1500*

available online at <http://research.ncl.ac.uk/nefg/QOF>

Systematic literature review: 343 peer-reviewed papers



17% studies were published before 2002, and 45% were published between 2008-2011


Data from supplementary Figure 1 in Baranski *et al.*, 2014

Systematic literature review: studies per crop type



Data from supplementary Table 3 in Baranski *et al.*, 2014

Primary assessment – antioxidant activity is significantly higher in organic crops



What are antioxidants/(poly)phenolics?

- **antioxidants neutralise free radicals**, thus preventing them from causing damage to cells in the body
- increased consumption of foods rich in antioxidants has been linked to a **reduction** in many **chronic diseases**, including:
 - cardiovascular disease
 - certain cancers (e.g. prostate cancer)
 - neurodegenerative diseases (Alzheimer's disease)
- plant-based (poly)phenolics are the most abundant antioxidants in the diet


Literature:

Del Rio D, Rodriguez-Mateos A, Spencer JPE *et al.* (2013) Dietary (poly)phenolics in human health: structures, bioavailability, and evidence of protective effects against chronic diseases. *Antioxid Redox Signal* 18, 1818-1892.

Wahlqvist ML (2013) Antioxidant relevance to human health. *Asia Pac J Clin Nutr* 22, 171-176.

Fardet A (2010) New hypotheses for the health-protective mechanisms of whole-grain cereals: what is beyond fibre? *Nutr Res Rev* 23, 65-134.

Explanatory assessments – antioxidant compounds



WHY do organic crops contain higher levels of antioxidants?

Evidence from the long-term, factorial experiments
at Nafferton Farm



NEFG Nafferton Farm

- 360 ha Estate
- cereals, oilseed
- dairy, beef, lamb
- sample preparation



Nafferton Factorial Systems Comparison Trial



experimental design - factors

- Rotation design (4)
 - Non-diverse (2)
 - 2 years grass/clover, 6 years cereals
 - 1 year potato or vegetables
 - Diversified (2)
 - 3 years grass/clover, 2 years cereals
 - 2 years potato or vegetables, 1 year faba beans
- Crop protection (2)
 - Conventional (based on pesticides)
 - Organic (according to soil association standards)
- Fertilisation (2)
 - Conventional (based on mineral fertilisers)
 - Organic (composted manure inputs only)
- Replicate blocks (4)
- Replicate experiments (4)
- Total area 6 ha


Nafferton Factorial Systems Comparison Trial experimental design - assessments

- Crop yield and quality parameters
- Disease, pest and weed incidence/pressure
- Gene, protein and metabolite expression
- Soil physical, chemical and biological parameters (including soil carbon sequestration)
- Environmental impact (e.g. N_2O , NO_3 and P losses) and life cycle analyses (LCA) for GHG emissions
- Resource use efficiency (e.g. fertiliser and crop protection inputs)
- Economic sustainability related parameters
- Biodiversity (e.g. non-crop plants and invertebrates)

Effect of fertilisation and crop protection on the glucosinolate content in cabbage (average of 2 seasons)



Primary assessment – cadmium concentration are significantly lower in organic crops



WHY do organic crops contain lower levels of cadmium?


Evidence from the long-term, factorial experiments
at Nafferton Farm




Effect of fertilisation and crop protection on the cadmium content in potato

(average of 4 seasons)


- mineral P-fertilisers contain cadmium (Cd)
- a range of scientific studies showed a close link between P-fertiliser use and cadmium concentrations in crops
- (Cooper *et al.*, 2011)



Primary assessment – pesticide residues are less frequently detected in organic crops



Explanatory assessment – pesticide residues in different crop types



Pesticide concentrations in organic and conventional crops

- There were too few studies/data-sets to compare pesticide concentrations in organic and conventional crops
- In the few individual studies where pesticide concentrations were compared, concentrations in conventional crops were 10 to 100 times higher

Why is the frequency of pesticide residues lower in organic crops?

- The use of synthetic chemical pesticides is prohibited in organic farming
- Organic farmers stick to the rules!!!

www.nefg-organic.org



Thank you